

DOROTA BRÓD

On a two-parameter generalization of Jacobsthal numbers and its graph interpretation

ABSTRACT. In this paper we introduce a two-parameter generalization of the classical Jacobsthal numbers ((s, p) -Jacobsthal numbers). We present some properties of the presented sequence, among others Binet's formula, Cassini's identity, the generating function. Moreover, we give a graph interpretation of (s, p) -Jacobsthal numbers, related to independence in graphs.

1. Introduction. The Jacobsthal sequence $\{J_n\}$ is defined by the second order linear recurrence

$$(1) \quad J_n = J_{n-1} + 2J_{n-2} \quad \text{for } n \geq 2$$

with $J_0 = 0$, $J_1 = 1$. The Binet's formula of this sequence has the following form

$$J_n = \frac{1}{3}(2^n - (-1)^n) \quad \text{for } n \geq 0.$$

Moreover, the explicit closed form expression for numbers J_n is

$$J_n = \sum_{r=0}^{\lfloor \frac{n-1}{2} \rfloor} \binom{n-1-r}{r} 2^r \quad \text{for } n \geq 0.$$

2010 *Mathematics Subject Classification.* 11B37, 11C20, 15B36, 05C69.

Key words and phrases. Jacobsthal numbers, generalized Jacobsthal numbers, Binet's formula, generating function, graph interpretation, Merrifield–Simmons index.

Other interesting properties of Jacobsthal numbers are given in [6]. There are many generalizations of this sequence in the literature. The second order recurrence (1) has been generalized in two ways: first, by preserving the initial conditions and second, by preserving the recurrence relation. We recall some of such generalizations:

- 1) k -Jacobsthal sequence $\{j_{k,n}\}$ [5], $j_{k,n+1} = kj_{k,n} + 2j_{k,n-1}$ for $k \geq 1$ and $n \geq 1$ with $j_{k,0} = 0$, $j_{k,1} = 1$,
- 2) k -Jacobsthal sequence $\{J_{k,n}\}$ [3], $J_{k,n+1} = J_{k,n} + kJ_{k,n-1}$ for $k \geq 1$ and $n \geq 1$ with $J_{k,0} = 0$, $J_{k,1} = 1$,
- 3) generalized Jacobsthal p -sequence $\{J_p\}$ [1], for any $p \in \mathbb{Z}^+$ and $n > p+1$ $J_p(n) = J_p(n-1) + 2J_p(n-p-1)$ with initial conditions $J_p(1) = J_p(2) = \dots = J_p(p+1) = 1$,
- 4) (s, t) -Jacobsthal sequence $\{\hat{j}_n(s, t)\}$ [8], $\hat{j}_n(s, t) = s\hat{j}_{n-1}(s, t) + 2t\hat{j}_{n-2}(s, t)$ for $n \geq 2$ with $\hat{j}_0(s, t) = 0$ and $\hat{j}_1(s, t) = 1$, for real numbers s, t , $s > 0$, $t \neq 0$ and $s^2 + 8t > 0$,
- 5) Jacobsthal sequence $\{J(d, t, n)\}$ [7], $J(d, t, n) = J(d, t, n-1) + tJ(d, t, n-d)$ for $n \geq d$ with $J(d, t, 0) = 1$, $J(d, t, n) = 1$ for $n = 1, \dots, d$, $t \geq 1$, $d \geq 2$.

In this paper we introduce a new generalization of the classical Jacobsthal numbers. Unlike other variations, this generalization depends on two integer parameters used in the recurrence relation (1). Let $n, s, p \geq 0$ be integers. We define (s, p) -Jacobsthal sequence $\{J_n(s, p)\}$ by the following recurrence

$$(2) \quad J_n(s, p) = 2^{s+p}J_{n-1}(s, p) + (2^{2s+p} + 2^{s+2p})J_{n-2}(s, p) \text{ for } n \geq 2$$

with initial conditions $J_0(s, p) = 1$, $J_1(s, p) = 2^s + 2^p + 2^{s+p}$.

For $s = p = 0$ we obtain $J_n(0, 0) = J_{n+2}$.

We will describe the terms of the sequence $\{J_n(s, p)\}$ explicitly by using a generalization of Binet's formula. Moreover, we will present some identities for (s, p) -Jacobsthal numbers, which generalize known results for the classical Jacobsthal numbers.

2. A graph interpretation of (s, p) -Jacobsthal numbers. In general we use the standard terminology and notation of graph theory, see [2]. In this section, we will present an interpretation of (s, p) -Jacobsthal numbers related to independence in graphs. Let G be a finite, undirected, simple graph with vertex set $V(G)$ and edge set $E(G)$. Recall that a subset S of $V(G)$ is an independent set of G if no two vertices of S are adjacent in G . Moreover, every one-element subset of $V(G)$ and the empty set are independent sets of G . The number of independent sets of a graph G is denoted by $NI(G)$. In the chemical literature the number of independent sets of a graph G is called the Merrifield–Simmons index of G and is denoted by $\sigma(G)$ ([4]). The numbers $J_n(s, p)$ have the graph interpretation directly related to the Merrifield–Simmons index.

Consider a graph $H_n^{s,p}$ (Figure 1), where $n \geq 1, s, p \geq 0$.




Figure 1.

Theorem 1. *Let n, s, p be integers, $n \geq 1, s, p \geq 0$. Then*

$$\sigma(H_n^{s,p}) = J_n(s, p).$$

Proof. In the beginning we will determine the number of independent sets of graphs $H_1^{s,p}$ and $H_2^{s,p}$. Assume that vertices of the graphs are numbered as in Figure 1. Denote by $L(x)$ the set of pendant vertices attached to the vertex x . Let $n = 1$. Assume that S is any independent set of $H_1^{s,p}$. Consider two cases.

Case 1. $y_1 \in S$.

Then $x_1, w_1, \dots, w_p \notin S$. Hence $S = \{y_1\} \cup Z$, where Z is any subset of the set $\{z_1, \dots, z_s\}$.

Case 2. $y_1 \notin S$. Consider two possibilities.

2.1. $x_1 \in S$.

Then $S = \{x_1\} \cup W$, where W is any subset of the set $\{w_1, \dots, w_p\}$.

2.2. $x_1 \notin S$.

Then $S = Z \cup W$.

Finally, we have $\sigma(H_1^{s,p}) = 2^s + 2^p + 2^{s+p} = J_1(s, p)$.

In the same manner we can obtain

$$\begin{aligned} \sigma(H_2^{s,p}) &= 2^{2p+s} + 2^{p+2s} + 2^{s+p}(2^s + 2^p + 2^{s+p}) \\ &= 2^{2s+p+1} + 2^{s+2p+1} + 2^{2s+2p} = J_2(s, p). \end{aligned}$$

Let $n \geq 3$. Assume that S is any independent set of $H_n^{s,p}$. Consider two cases.

Case 1. $y_n \in S$.

Let \mathcal{S}_1 be a family of all independent sets S of the graph $H_n^{s,p}$ such that $y_n \in S$. Then $x_n, x_{n-1}, y_{n-1}, w_1, \dots, w_p \notin S$. Hence $S = S' \cup \{y_n\} \cup S_1 \cup S_2 \cup S_3$, where S' is any independent set of the graph $H_n^{s,p} \setminus \{x_n, x_{n-1}, y_n, y_{n-1}\} \setminus (L(x_n) \cup L(x_{n-1}) \cup L(y_n) \cup L(y_{n-1}))$, isomorphic to $H_{n-2}^{s,p}$, $S_1 \subset L(x_n)$, $S_2 \subset L(x_{n-1})$, $S_3 \subset L(y_{n-1})$. Hence by the fundamental combinatorial statements we have $|\mathcal{S}_1| = 2^p \cdot (2^s)^2 \sigma(H_{n-2}^{s,p})$.

Case 2. $y_n \notin S$.

Let \mathcal{S}_2 be a family of all independent sets S of the graph $H_n^{s,p}$ such that $y_n \notin S$. Consider two possibilities.

2.1. $x_n \notin S$.

Then $S = S'' \cup S_1 \cup S_4$, where S'' is any independent set of the graph $H_n^{s,p} \setminus \{x_n, y_n\} \setminus (L(x_n) \cup L(y_n))$, isomorphic to $H_{n-1}^{s,p}$, $S_1 \subset L(x_n)$, $S_4 \subset L(y_n)$.

2.2. $x_n \in S$.

Then $S = S' \cup \{x_n\} \cup S_2 \cup S_3 \cup S_4$, where S' is any independent set of the graph $H_n^{s,p} \setminus \{x_n, x_{n-1}, y_n, y_{n-1}\} \setminus (L(x_n) \cup L(x_{n-1}) \cup L(y_n) \cup L(y_{n-1}))$, isomorphic to $H_{n-2}^{s,p}$.

Consequently, $|\mathcal{S}_2| = 2^s \cdot 2^p \sigma(H_{n-1}^{s,p}) + (2^p)^2 \cdot 2^s \sigma(H_{n-2}^{s,p})$.

Finally, for $n \geq 3$ we obtain

$$\sigma(H_n^{s,p}) = |\mathcal{S}_1| + |\mathcal{S}_2| = 2^{s+p} \sigma(H_{n-1}^{s,p}) + (2^{2s+p} + 2^{2p+s}) \sigma(H_{n-2}^{s,p})$$

with $\sigma(H_1^{s,p}) = 2^s + 2^p + 2^{s+p}$ and $\sigma(H_2^{s,p}) = 2^{2s+p+1} + 2^{s+2p+1} + 2^{2s+2p}$, which ends the proof. \square

Corollary 2. *Let $n \geq 1$. Then $\sigma(H_n^{0,0}) = J_n(0,0) = J_{n+2}$.*

3. Some identities for (s, p) -Jacobsthal numbers. The characteristic equation, associated with the recurrence relation (2) is

$$(3) \quad r^2 - 2^{s+p}r - (2^{2s+p} + 2^{s+2p}) = 0$$

with roots

$$(4) \quad r_1 = 2^{s+p-1} + \frac{1}{2} \sqrt{4^{s+p} + 2^{s+p+2}(2^s + 2^p)},$$

$$(5) \quad r_2 = 2^{s+p-1} - \frac{1}{2} \sqrt{4^{s+p} + 2^{s+p+2}(2^s + 2^p)}.$$

Note that

$$(6) \quad r_1 + r_2 = 2^{s+p},$$

$$(7) \quad r_1 r_2 = -(2^{2s+p} + 2^{s+2p}),$$

$$(8) \quad r_1 - r_2 = \sqrt{4^{s+p} + 2^{s+p+2}(2^s + 2^p)}.$$

The general formula of (s, p) -Jacobsthal sequence can be written by the following identity

$$J_n(s, p) = c_1 r_1^n + c_2 r_2^n$$

for some constants c_1, c_2 . Using initial conditions $J_0(s, p) = 1$, $J_1(s, p) = 2^s + 2^p + 2^{s+p}$, we get the system of two linear equations

$$\begin{cases} c_1 + c_2 = 1 \\ c_1 r_1 + c_2 r_2 = 2^s + 2^p + 2^{s+p}. \end{cases}$$

Solving the system, we obtain

$$(9) \quad \begin{aligned} c_1 &= \frac{2^s + 2^p + 2^{s+p} - 2^{s+p-1} + \frac{1}{2}\sqrt{4^{s+p} + 2^{s+p+2}(2^s + 2^p)}}{\sqrt{4^{s+p} + 2^{s+p+2}(2^s + 2^p)}}, \\ c_2 &= \frac{2^{s+p-1} - 2^s - 2^p - 2^{s+p} + \frac{1}{2}\sqrt{4^{s+p} + 2^{s+p+2}(2^s + 2^p)}}{\sqrt{4^{s+p} + 2^{s+p+2}(2^s + 2^p)}}. \end{aligned}$$

Hence we get the following result.

Proposition 3 (Binet's formula). *Let $n, s, p \geq 0$. Then the n -th (s, p) -Jacobsthal number is given by*

$$(10) \quad \begin{aligned} J_n(s, p) &= \frac{(2^s + 2^p + 2^{s+p} - 2^{s+p-1} + \frac{1}{2}\sqrt{\Delta})r_1^n}{\sqrt{\Delta}} \\ &\quad + \frac{(2^{s+p-1} - 2^s - 2^p - 2^{s+p} + \frac{1}{2}\sqrt{\Delta})r_2^n}{\sqrt{\Delta}}, \end{aligned}$$

where $\Delta = 4^{s+p} + 2^{s+p+2}(2^s + 2^p)$, $r_1 = 2^{s+p-1} + \frac{1}{2}\sqrt{\Delta}$, $r_2 = 2^{s+p-1} - \frac{1}{2}\sqrt{\Delta}$.

Using Binet's formula, we can get some identities for (s, p) -Jacobsthal numbers.

Theorem 4 (Cassini's identity). *Let n, s, p be integers, $n \geq 1$, $s, p \geq 0$. Then*

$$J_{n+1}(s, p)J_{n-1}(s, p) - J_n^2(s, p) = (-1)^n(2^s + 2^p)^2(2^{2s+p} + 2^{s+2p})^{n-1}.$$

Proof. By formula (10) we get

$$\begin{aligned} &J_{n+1}(s, p)J_{n-1}(s, p) - J_n^2(s, p) \\ &= (c_1r_1^{n+1} + c_2r_2^{n+1})(c_1r_1^{n-1} + c_2r_2^{n-1}) - (c_1r_1^n + c_2r_2^n)^2 \\ &= c_1c_2r_1^{n+1}r_2^{n-1} + c_1c_2r_2^{n+1}r_1^{n-1} - 2c_1c_2r_1^n r_2^n \\ &= c_1c_2(r_1r_2)^n \left(\frac{r_1}{r_2} + \frac{r_2}{r_1} - 2\right) = c_1c_2(r_1r_2)^{n-1}(r_1 - r_2)^2. \end{aligned}$$

By simple calculations we obtain

$$(11) \quad c_1c_2 = \frac{-(2^s + 2^p)^2}{4^{s+p} + 2^{s+p+2}(2^s + 2^p)}.$$

Using formulas (7), (8) and (11), we have

$$J_{n+1}(s, p)J_{n-1}(s, p) - J_n^2(s, p) = (-1)^n(2^s + 2^p)^2(2^{2s+p} + 2^{s+2p})^{n-1}.$$

□

Proposition 5. *Let n, s, p be integers, $n \geq 1$, $s, p \geq 0$. Then*

$$\lim_{n \rightarrow \infty} \frac{J_{n+1}(s, p)}{J_n(s, p)} = 2^{s+p-1} + \frac{1}{2}\sqrt{4^{s+p} + 2^{s+p+2}(2^s + 2^p)}.$$

Proof. By formula (10) we have

$$\lim_{n \rightarrow \infty} \frac{J_{n+1}(s, p)}{J_n(s, p)} = \lim_{n \rightarrow \infty} \frac{c_1 r_1^{n+1} + c_2 r_2^{n+1}}{c_1 r_1^n + c_2 r_2^n} = \lim_{n \rightarrow \infty} \frac{c_1 r_1 + c_2 r_2 \left(\frac{r_2}{r_1}\right)^n}{c_1 + c_2 \left(\frac{r_2}{r_1}\right)^n}.$$

Since $\lim_{n \rightarrow \infty} \left(\frac{r_2}{r_1}\right)^n = 0$, we have

$$\lim_{n \rightarrow \infty} \frac{J_{n+1}(s, p)}{J_n(s, p)} = r_1 = 2^{s+p-1} + \frac{1}{2} \sqrt{4^{s+p} + 2^{s+p+2}(2^s + 2^p)}.$$

□

Theorem 6 (summation formula).

$$(12) \quad \sum_{i=0}^{n-1} J_i(s, p) = \frac{J_n(s, p) + (2^{2s+p} + 2^{s+2p})J_{n-1}(s, p) - 1 - 2^s - 2^p}{2^{s+p}(1 + 2^s + 2^p) - 1}.$$

Proof. By Binet's formula (10) we have

$$\begin{aligned} \sum_{i=0}^{n-1} J_i(s, p) &= \sum_{i=0}^{n-1} (c_1 r_1^i + c_2 r_2^i) = c_1 \frac{1 - r_1^n}{1 - r_1} + c_2 \frac{1 - r_2^n}{1 - r_2} \\ &= \frac{c_1 + c_2 - (c_1 r_2 + c_2 r_1) - (c_1 r_1^n + c_2 r_2^n) + r_1 r_2 (c_1 r_1^{n-1} + c_2 r_2^{n-1})}{(1 - r_1)(1 - r_2)} \\ &= \frac{c_1 + c_2 - (c_1 r_2 + c_2 r_1) - J_n(s, p) + r_1 r_2 J_{n-1}(s, p)}{1 - (r_1 + r_2) + r_1 r_2}. \end{aligned}$$

By formulas (4), (5) and (9) we obtain

$$(13) \quad c_1 r_2 + c_2 r_1 = -(2^s + 2^p).$$

Using (6), (7) and (13), we get

$$\sum_{i=0}^{n-1} J_i(s, p) = \frac{1 + 2^s + 2^p - J_n(s, p) - (2^{2s+p} + 2^{s+2p})J_{n-1}(s, p)}{1 - 2^{s+p} - 2^{2s+p} - 2^{s+2p}}.$$

Hence

$$\sum_{i=0}^{n-1} J_i(s, p) = \frac{J_n(s, p) + (2^{2s+p} + 2^{s+2p})J_{n-1}(s, p) - 1 - 2^s - 2^p}{2^{s+p}(1 + 2^s + 2^p) - 1}.$$

□

Corollary 7. For $s = p = 0$ we get the well-known identity for the classical Jacobsthal numbers

$$\sum_{i=0}^{n-1} J_i = \frac{J_{n+2} + 2J_{n+1} - 3}{2}.$$

The next theorem presents the generating function of (s, p) -Jacobsthal sequence.

Theorem 8. *The generating function of the sequence $\{J_n(s, p)\}$ has the following form*

$$f(x) = \frac{1 + (2^s + 2^p)x}{1 - 2^{s+p}x - (2^{2s+p} + 2^{s+2p})x^2}.$$

Proof. Let $f(x) = \sum_{n=0}^{\infty} J_n(s, p)x^n$. Then, by recurrence relation (2), we have

$$\begin{aligned} f(x) &= J_0(s, p) + J_1(s, p)x + \sum_{n=2}^{\infty} J_n(s, p)x^n \\ &= 1 + (2^s + 2^p + 2^{s+p})x \\ &\quad + \sum_{n=2}^{\infty} (2^{s+p}J_{n-1}(s, p) + (2^{2s+p} + 2^{s+2p})J_{n-2}(s, p))x^n \\ &= 1 + (2^s + 2^p + 2^{s+p})x \\ &\quad + 2^{s+p} \sum_{n=2}^{\infty} J_{n-1}(s, p)x^n + (2^{2s+p} + 2^{s+2p}) \sum_{n=2}^{\infty} J_{n-2}(s, p)x^n \\ &= 1 + (2^s + 2^p + 2^{s+p})x \\ &\quad + 2^{s+p}x \sum_{n=1}^{\infty} J_n(s, p)x^n + (2^{2s+p} + 2^{s+2p})x^2 \sum_{n=0}^{\infty} J_n(s, p)x^n \\ &= 1 + (2^s + 2^p + 2^{s+p})x \\ &\quad + 2^{s+p}x \sum_{n=0}^{\infty} J_n(s, p)x^n - 2^{s+p}x + (2^{2s+p} + 2^{s+2p})x^2 f(x). \end{aligned}$$

Thus

$$f(x) = 1 + (2^s + 2^p)x + 2^{s+p}xf(x) + (2^{2s+p} + 2^{s+2p})x^2f(x).$$

Hence

$$f(x) = \frac{1 + (2^s + 2^p)x}{1 - 2^{s+p}x - (2^{2s+p} + 2^{s+2p})x^2},$$

which ends the proof. □

REFERENCES

- [1] Dasdemir, A., *The representation, generalized Binet formula and sums of the generalized Jacobsthal p-sequence*, Hittite Journal of Science and Engineering **3** (2) (2016), 99–104.
- [2] Diestel, R., *Graph Theory*, Springer-Verlag, Heidelberg–New York, 2005.
- [3] Falcon, S., *On the k-Jacobsthal numbers*, American Review of Mathematics and Statistics **2** (1) (2014), 67–77.

- [4] Gutman, I., Wagner, S., *Maxima and minima of the Hosoya index and the Merrifield-Simmons index: a survey of results and techniques*, Acta Appl. Math. **112** (3) (2010), 323–348.
- [5] Jhala, D., Sisodiya, K., Rathore, G. P. S., *On some identities for k -Jacobsthal numbers*, Int. J. Math. Anal. (Ruse) **7** (9–12) (2013), 551–556.
- [6] Horadam, A. F., *Jacobsthal representation numbers*, Fibonacci Quart. **34** (1) (1996), 40–54.
- [7] Szynal-Liana, A., Włoch, A., Włoch, I., *On generalized Pell numbers generated by Fibonacci and Lucas numbers*, Ars Combin. **115** (2014), 411–423.
- [8] Uygun, S., *The (s, t) -Jacobsthal and (s, t) -Jacobsthal Lucas sequences*, Applied Mathematical Sciences **9** (70) (2015), 3467–3476.

Dorota Bród
Rzeszów University of Technology
Faculty of Mathematics and Applied Physics
al. Powstańców Warszawy 12
35-959 Rzeszów
Poland
e-mail: dorotab@prz.edu.pl

Received March 1, 2018